$SU(3)$ colorsingletness, $Z(3)$ symmetry, Polyakov Loop and dynamical recombination

Chowdhury Aminul Islam[a], Raktim Abir[a], Munshi G. Mustafa[a], Sanjay K. Ghosh[b] and Rajarshi Ray[b]
[a] *Theory Division, Saha Institute of Nuclear Physics,*
1/AF Bidhan Nagar, Kolkata-700 064, INDIA and
[b] *Center for Astroparticle Physics and Space Science,*
Bose Institute, 5/EN Bidhan Nagar, Kolkata-700091, INDIA &
Department of Physics, Bose Institute, 93/1, A. P. C Road, Kolkata - 700009, INDIA
(Dated: August 16, 2012)

arXiv:1208.3146v1 [hep-ph] 15 Aug 2012

Based on quantum statistical mechanics we show that the $SU(3)$ colorsinglet ensemble of a quark-gluon gas exhibits a $Z(3)$ symmetry through the normaized character in fundamental representation and also becomes equivalent, within a stationary point approximation, to the ensemble given by Polyakov Loop. The probability of the normalized character in $SU(3)$ is found to be maximum at a particular value exhibiting a long range color correlation. This clearly indicates a transition from a color correlated to uncorrelated phase or vise-versa. A dynamical recombination of ionized $Z(3)$ color charges to a color singlet $Z(3)$ symmetric confined phase is evident along with a lower bound that originates from an exchange of a pair of massive magnetic gluons between two Polyakov Loops.

PACS numbers: 12.38.Mh, 25.75.+r, 24.85.+p, 25.75.-q, 25.75.Nq

Introduction: A statistical thermodynamic description of a quantum gas, is often useful for various physical systems, *e.g.*, electrons in metal, blackbody photons in a heated cavity, phonons at low temperature, neutron matter in neutron stars, etc. In most cases the mutual interactions among the constituents are neglected, although they interact in order to come to a thermal equilibrium. One can imagine a situation by first allowing them to come to a thermal equilibrium and then slowly turning off the interactions [1]. In the same perspective we can consider a quantum gas of non-interacting quarks (q), antiquarks ($\bar{q}$) and gluons (g) with the underlying symmetry of $SU(3)$ color gauge theory at a given temperature. Such a simple quantum statistical description exhibits very interesting features of a quark-gluon system.

Quantum statistical mechanics and colorsinglet ensemble: In thermal equilibrium the statistical behaviour of a quantum gas is studied through a density matrix in an appropriate ensemble in which the partition function in a finite volume is defined as

$$\mathcal{Z} = \mathrm{Tr}\left(\hat{\mathcal{P}} e^{-\beta \hat{H}}\right) = \sum_n \left\langle n \left| \hat{\mathcal{P}} e^{-\beta \hat{H}} \right| n \right\rangle \quad , \qquad (1)$$

where $|n\rangle$ is a many-particle state, β is the inverse of temperature T and $\hat{H}$ is the Hamiltonian. Let $\mathcal{G}$ be a symmetry group with unitary representation $\hat{U}(g)$ in a Hilbert space $\mathcal{H}$. The projection operator [2–6] $\hat{\mathcal{P}}$ for a desired configuration is defined as $\hat{\mathcal{P}}_j = d_j \int_{\mathcal{G}} \mathrm{d}\mu(g) \chi_j^\star(g) \hat{U}(g)$, where d_j and χ_j are, respectively, the dimension and the character of the irreducible representation j of $\mathcal{G}$ and $\mathrm{d}\mu(g)$ is the invariant Haar measure. The symmetry group associated with the color singlet configuration is $SU(N_c)$ and $d_j = 1$ and $\chi_j = 1$. Now the color singlet partition function for the system becomes,

$$\mathcal{Z}_S = \int_{SU(N_c)} \mathrm{d}\mu(g) \mathrm{Tr}\left(\hat{U}(g) \exp(-\beta \hat{H})\right) \quad . \qquad (2)$$

The invariant Haar measure [2, 3] is expressed in terms of the distribution of eigenvalues of $SU(N_c)$ as

$$\int \mathrm{d}\mu(g) = \frac{1}{N_c!} \left(\prod_{i=1}^{N_c} \int_{-\pi}^{\pi} \frac{\mathrm{d}\theta_l}{2\pi} \right) \delta\Big(\sum_i \theta_i\Big) \prod_{i>j} \left| e^{i\theta_i} - e^{i\theta_j} \right|^2 \quad (3)$$

where the product of the differences of the eigenvalues is known as the Vandermonde (VdM) determinant. The class parameter θ_l obeys $\sum_{l=1}^{N_c} \theta_l = 0 \ (\mathrm{mod} 2\pi)$ ensuring the requirement of unit determinant in $SU(N_c)$. This also restricts that $SU(3)$ has only two parameter abelian subgroups associated with two diagonal generators, which would completely characterize the $\hat{U}(g)$.

Now, the Hilbert space $\mathcal{H}$ of a composite system has a structure of a tensor product of the individual Fock spaces as $\mathcal{H} = \mathcal{H}_q \otimes \mathcal{H}_{\bar{q}} \otimes \mathcal{H}_g$. The partition function in (2) decomposes [3–6] in respective Fock spaces as,

$$\mathcal{Z}_S = \int_{SU(N_c)} \mathrm{d}\mu(g) \mathrm{Tr}\left(\hat{U}_q e^{-\beta \hat{H}_q}\right) \mathrm{Tr}\left(\hat{U}_{\bar{q}} e^{-\beta \hat{H}_{\bar{q}}}\right) \mathrm{Tr}\left(\hat{U}_g e^{-\beta \hat{H}_g}\right) , (4)$$

where the various $U_i(g)$ act as link variables that link, respectively, the quarks, antiquarks and spatial gluons in a given state of the physical system. In each Fock space there exists a basis that diagonalizes both operators as long as $\hat{H}_i$ and $\hat{U}_i$ commute. Performing the traces in (4) using the standard procedure [1, 3, 5], the partition function in Hilbert space becomes,

$$\mathcal{Z}_S = \int_{SU(N_c)} \mathrm{d}\mu(g) \;\; e^{\Theta_q + \Theta_{\bar{q}} + \Theta_g} = \int_{SU(N_c)} \mathrm{d}\mu(g) \;\; e^{\Theta} \quad ; \quad (5)$$

$$\begin{aligned} \Theta \; = \; & 2N_f \sum_\alpha \mathrm{tr_c} \ln \left(1 + R_q e^{-\beta(\epsilon_q^\alpha - \mu_q)}\right) \\ & + 2N_f \sum_\alpha \mathrm{tr_c} \ln \left(1 + R_{\bar{q}} e^{-\beta(\epsilon_q^\alpha + \mu_q)}\right) \\ & - 2 \sum_\alpha \mathrm{tr_c} \ln \left(1 - R_g e^{-\beta \epsilon_g^\alpha}\right) , \end{aligned} \qquad (6)$$



FIG. 1: (color online) A 3D-plot of $|\Phi(\theta_1, \theta_2)|$, i.e., the normalized character in the fundamental representation of $SU(3)$ as given in (8) within $-\pi \leq \theta_1,\, \theta_2 \leq \pi$.

where $\epsilon_i^\alpha = \sqrt{(p_i^\alpha)^2 + m_i^2}$. Also the flavor (N_f), spin and the chemical potential μ are introduced. The finite dimensional diagonal matrix $R_{q(\bar{q})}$ in the basis of the color space represents the image [2, 3] of the group element in the irreducible representation of $SU(N_c)$ as,

$$R_q \;=\; \mathrm{diag}\left(e^{i\theta_1}, e^{i\theta_2}, e^{i\theta_3}\right) \;;\; R_{\bar{q}} = R_q^\dagger. \qquad (7)$$

We also define normalised characters by the respective dimension of the fundmaental and adjoint representations as,

$$\Phi = \frac{1}{N_c}\mathrm{tr}_c R_q \;, \qquad \bar{\Phi} = \frac{1}{N_c}\mathrm{tr}_c R_q^\dagger \;,$$
$$(N_c^2 - 1)\Phi_A \;=\; N_c^2 \Phi\bar{\Phi} - 1\,. \qquad (8)$$

In Fig. 1 a three dimensional view of the magnitude of the normalized character in fundamental representation of $SU(3)$, $|\Phi(\theta_1, \theta_2)|$, is shown within the domain $-\pi \leq \theta_1, \theta_2 \leq \pi$. It has three maxima for $(\theta_1, \theta_2) = (0, 0)$, $(2\pi/3, 2\pi/3)$ and $(-2\pi/3, -2\pi/3)$. There are also three minima (and three mirror images exist if one interchanges $\theta_1 \leftrightarrow \theta_2$) for $(\theta_1, \theta_2) = (0, 2\pi/3)$, $(0, -2\pi/3)$ and $(2\pi/3, -2\pi/3)$. These three minima are the three vacua of Φ in (θ_1, θ_2)-plane and separated from each other by potential walls. $|\Phi|$ has a three fold degeneracy connected by a rotation of $2\pi/3$ in both θ_1 and θ_2. A Monte Carlo simulation of complex Φ is also displayed in Fig. 2 in a Argrand plane with $-\pi \leq (\theta_1,\, \theta_2) \leq \pi$. This shows a three pointed star in a circle of unit radius, in which each point can be rotated by a phase $2\pi/3$ except the origin. This clearly indicates that Φ in the fundamental representation of $SU(3)$ has a center symmetry $Z(3)$ with three rotational angles (viz., 0, $2\pi/3$, $4\pi/3$). This can also be understood from the invariant Haar measure expressed in terms of Φ and $\bar{\Phi}$. Now, the three minima in Fig. 1 uniquely correspond to the center of the circle at $\Phi = 0$ in Fig. 2, which is the $Z(3)$ symmetric phase or confined phase at low T. On the other hand, the three maxima in Fig. 1 correspond to the three pointed tips in Fig. 2 representing the spontaneously broken phase or deconfined phase of $Z(3)$ at very high T. Φ can thus act as an order parameter for deconfinement phase transition. In Fig. 3 a three dimensional plot of Φ_A is also displayed that exhibits same features as Fig. 1 except that the minima appear with negative values, which could be understood from (8).



FIG. 2: (color online) A Monte Carlo simulation of complex $\Phi(\theta_1, \theta_2)$ in Argrand plane for which θ_1 and θ_2 are chosen randomly in the domain $-\pi \leq \theta \leq \pi$.



FIG. 3: (color online) Same as Fig. 1 but for $\Phi_A(\theta_1, \theta_2)$.

The VdM term in (3) can now be written in terms of Φ and $\bar{\Phi}$ as $\prod_{i>j}^{3} \left|e^{i\theta_i} - e^{i\theta_j}\right|^2 = 27[1 - 6\Phi\bar{\Phi} + 4(\Phi^3 + \bar{\Phi}^3) - 3(\Phi\bar{\Phi})^2] = 27\ H(\Phi, \bar{\Phi})$, for $SU(3)$. Further, this is in general not possible for $N_c > 3$ as there are more than two independent parameters. Now the Jacobian for variable transformation from $\{\theta_1, \theta_2\}$ to $\{\Phi, \bar{\Phi}\}$ can be obtained as $J(\Phi, \bar{\Phi}) = (1/9)\sqrt{27H(\Phi, \bar{\Phi})}$. In the infinite volume V, one also needs to replace the discrete single particle sum by an integral as $\sum_\alpha \rightarrow (V/(2\pi)^3 \int d^3p$.

After performing the color trace of the matter and gauge parts, and expressing in terms of the characters of the fundamental and its conjugate, Eq.(6) becomes

$$\Theta = \Theta_q + \Theta_{\bar{q}} + \Theta_g = 2N_f V \int \frac{d^3p}{(2\pi)^3} \ln\Big[1 + e^{-3\beta\epsilon_q^+}$$
$$+ N_c\left(\Phi + \bar{\Phi}e^{-\beta\epsilon_q^+}\right)e^{-\beta\epsilon_q^+}\Big] + 2N_f V \int \frac{d^3p}{(2\pi)^3} \ln\Big[1 +$$
$$e^{-3\beta\epsilon_q^-} + N_c\left(\bar{\Phi} + \Phi e^{-\beta\epsilon_q^-}\right)e^{-\beta\epsilon_q^-}\Big]$$
$$-2TV \int \frac{d^3p}{(2\pi)^3} \ln\left(1 + \sum_{n=1}^{8} a_n\, e^{-n\beta\epsilon_g}\right). \qquad (9)$$

with $\epsilon_q^\pm = \epsilon_q \mp \mu$ and the coefficients a_n are given by

$$\begin{aligned}
a_1 &= a_7 = 1 - N_c^2\bar{\Phi}\Phi\,, \quad a_8 = 1\,,\\
a_2 &= a_6 = 1 - 3N_c^2\bar{\Phi}\Phi + N_c^3\left(\bar{\Phi}^3 + \Phi^3\right)\,,\\
a_4 &= 2\left[-1 + N_c^2\bar{\Phi}\Phi - N_c^3\left(\bar{\Phi}^3 + \Phi^3\right) + N_c^4\left(\bar{\Phi}\Phi\right)^2\right]\,,\\
a_3 &= a_5 = -2 + 3N_c^2\bar{\Phi}\Phi - N_c^4\left(\bar{\Phi}\Phi\right)^2\,. \qquad (10)
\end{aligned}$$

Now in the infinite volume limit the color singlet partition function in (5)

$$\mathcal{Z}_S \;=\; \int_{SU(3)} d\Phi\; d\bar{\Phi}\; e^{\Theta_q+\Theta_{\bar{q}}+\Theta_g+\Theta_H}, \qquad (11)$$

with $\Theta_H = \frac{1}{2}\ln H + \ln\frac{9\sqrt{3}}{8\pi^2} \approx \frac{1}{2}\ln H$, where the constant normalization factor is dropped as it is subleading.

Now performing the integrations using the method of stationary points, one can write

$$\mathcal{Z}^0_S = e^{\Theta_q+\Theta_{\bar{q}}+\Theta_g+\Theta_{\rm H}}\Big|_{\substack{\Phi\to\Phi_0\\ \bar{\Phi}\to\bar{\Phi}_0}} = \mathcal{Z}^{\rm PL}(\Phi_0,\bar{\Phi}_0)\,, \qquad (12)$$

where 'PL' stands for Polyakov Loop [7]. This exhibits that the $SU(3)$ color singlet ensemble of a quark-gluon gas is equivalent to that of the Polyakov Loop at the stationary values of Φ_0 and $\bar{\Phi}_0$, which can be obtained from the extremum conditions as

$$\frac{\partial}{\partial\Phi}\left[\Theta+\Theta_{\rm H}\right] \;=\; 0\,;\; \frac{\partial}{\partial\bar{\Phi}}\left[\Theta+\Theta_{\rm H}\right] = 0\,. \qquad (13)$$

The Wilson line in $SU(N_c)$ is defined [7] as $L = \mathcal{T}\exp[ig\int\limits_0^\beta A_0\; dt]$, where A_0 is a temporal gauge field, g is the strong coupling constant and $\mathcal{T}$ is the path-ordering in Euclidean time. The trace of the Wilson line, known as Polyakov Loop, is an order parameter for pure gauge theories. The gauge invariant Polyakov Loop fields may be expressed in terms of the characters of the representation as given in (8).

The color singlet thermodynamic potential density at the stationary points in infinite volume limit becomes

$$\begin{aligned}\Omega_S \;&=\; -\frac{T}{V}\ln\mathcal{Z}^0_S = -\frac{T}{V}\left[\Theta_q+\Theta_{\bar{q}}+\Theta_g+\Theta_{\rm H}\right]_{\substack{\Phi\to\Phi_0\\ \bar{\Phi}\to\bar{\Phi}_0}}\\ &= \Big[\Omega_q+\Omega_{\bar{q}}+\Omega_g+\Omega_{\rm H}\Big]_{\substack{\Phi\to\Phi_0\\ \bar{\Phi}\to\bar{\Phi}_0}} = \Omega^{\rm PL}(\Phi_0,\bar{\Phi}_0). \qquad (14)\end{aligned}$$

As discussed the normalized character in fundamental representation vis-a-vis PL transforms under the global $Z(N_c)$ symmetry as a field with charge one as $\Phi \rightarrow e^{2\pi i/N_c}\Phi$. Given the role of an order parameter for pure gauge [7], if $\Phi = 0$ the $Z(N_c)$ is unbroken and there is no ionization of $Z(N_c)$ charge, in the confined phase below a certain temperature. At high temperature the symmetry is spontaneously broken, $\Phi \neq 0$ corresponds to a deconfined phase of gluonic plasma and there are N_c different equilibrium states distinguished by the phase $2\pi j/N_c$ with $j = 0, \cdots N_c - 1$ as shown in Figs. 1 and 2. Both deconfinement [7] and chiral dynamics [10] have been studied extensively using PNJL model [8] in which PL with mostly (anti)quarks as quasiparticles is usually coupled to NJL model [9]. In some cases only the gluons as quasiparticles in PL model was also studied [11, 12]. Here we have derived the PL model phenomenologically by imposing $SU(N_c)$ color singlet restriction on a quark-gluon gas, which could easily be coupled to any chiral model to study deconfinement and chiral dynamics.



FIG. 4: (color online) A Monte Carlo simulation, $P(\Phi)$, corresponds to the occurrence probability of $|\Phi(\theta_1,\theta_2)|$ in $SU(3)$ parameter space $(\theta_1,\, \theta_2)$. The values of θ_1 and θ_2 are chosen randomly within the domain $-\pi \leq \theta_1,\, \theta_2 \leq \pi$ and then the obtained value of $|\Phi|$ is mapped in a given Φ bin, which is then normalized by the area of the bin.

However, we also noticed some more interesting features of Φ in $(\theta_1,\, \theta_2)$-plane. In Fig. 4 a Monte Carlo simulation of the occurrence probability, $P(|\Phi|)$, of $|\Phi(\theta_1,\theta_2)|$ is displayed in $SU(3)$ parameter space. This plot indicates that the most probable value for $|\Phi|$ is 1/3. As T increases, Φ increases [8] from zero in the confined phase and reaches unity for ideal gas. In the domain $0 < |\Phi| < 1/3$, the color states begin to show up from the color neutral states which have not yet melted completely but prefer to reside in $Z(3)$ vacua. This domain is shown by red dots ($|\Phi| \sim 0$) to purple triangles ($|\Phi| \sim 0.3$) in Fig. 5, which is a contour plot corresponding to Fig. 1. As long as such states are inside the domain of $Z(3)$ vacua, a strong localized color correlation exists among the color charges like a liquid. In this $|\Phi|$ domain, the normalized character in adjoint representation varies as $-1/8 \leq \Phi_A < 0$ which is represented in Fig. 6, which is again a contour plot corresponding to Fig. 3. Now for $|\Phi| = 1/3$, the color charges begin to disperse from the $Z(3)$ vacua and get connected to all other color charges in $Z(3)$ domain, which is represented by the yellow mesh in Fig. 5. This causes $P|\Phi|$ to be maximum in Fig. 4 exhibiting a long range color correlation. Here $\Phi_A = 0$ as is also represented by the blue mesh in Fig. 6. When $1/3 < |\Phi| \leq 1$, the color charges begin to get uncorrelated from each other and the ionization of $Z(3)$ color charges take place as obvious from equivalued $|\Phi|$ lines in Fig. 5 starting from sea-blue ($|\Phi| \sim 0.4$) to green lines ($|\Phi| \sim 0.9$) at a step of 0.1. When $|\Phi| \sim 1$ a complete ionization of $Z(3)$ charges take place and they reside at those maxima in Fig. 1 which are also represented by blue blobs in Fig. 5. This ionization can also be seen in Fig. 6 through purple equivalued Φ_A lines to purple blobs in the range $0.1 \leq \Phi_A \leq 1$. All these features



FIG. 5: (color online) A 2D projection of $|\Phi|$ in θ_1 and θ_2 plane in which each color corresponds to a equivalued $|\Phi|$. The red dots to purple triangles correspond to $|\Phi| \sim 0, 0.1, 0.2, 0.3$ whereas sea-blue lines to blue dots correspond to $|\Phi| \sim 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1$. The equivalued mesh connected by yellow triangles correspond to $|\Phi| \sim 1/3$.



FIG. 6: (color online) Same as Fig. 5 but for Φ_A. The equivalued mesh connected by blue triangles is for $\Phi_A = 0$. From the blue mesh to purple blobs, Φ_A increases by a step of 0.1. The red dots are for $\Phi_A \sim -1/8$ whereas the green triangles $\Phi_A \sim -0.05$.

are also reflected in complex Φ plane in Fig. 7 indicating that three ionized color charges get recombined together and transformed into a color neutral state through $Z(3)$ vacua and vise-versa. A clear indication of a phase transition is evident in $SU(3)$ parameter space in the domain $0.3 \leq |\Phi| \leq 0.35$ implying a smooth transition from a correlated color states (strongly correlated quark-gluon plasma (QGP) like a liquid) to uncorrelated color states (QGP) at temperature T_c. Also T_s is a temperature corresponding to $|\Phi| \sim 0$ at which color neutral states begin to transform to a highly color localized phase or vise-versa. Moreover, the domain, $0 \leq |\Phi| \leq 1/3$ also exhibits a $U(1)$ symmetry in $Z(3)$ (Fig. 7). This domain is a metastable state with temperature range $T_s \leq T \leq T_c$, that indicates melting of color neutral state to highly color localized phase whereas the color ionization domain is $1/3 \leq |\Phi| \leq 1$ at $T > T_c$. All these features are also observed in a recent Lattice QCD calculation [13].

Matter sector: The $Z(N_c)$ symmetry in the thermodynamic potential for (anti)quarks, $\Omega_{q(\bar{q})}^{\rm PL}$, in (14) is explicitly broken under the rotation of $Z(N_c)$ since they also carry the $Z(N_c)$ charge. In the asymptotically high temperature ($T \gg T_c$) the matter part in (14) becomes



FIG. 7: (color online) The projection of Φ in Argrand plane for a fixed value, which are noted in the respective plots. This shows a strong color correlation in the range $0 \leq \Phi \leq 1/3$ and ionization of color states in the range $1/3 < \Phi \leq 1$.

$$\Omega_{q(\bar{q})}^{\Phi_0,\bar{\Phi}_0\to 1} = -2N_f T\int\frac{d^3p}{(2\pi)^3}\ln\left(1+e^{-\beta(\epsilon_q\mp\mu)}\right)^{N_c} \quad (15)$$

where $N_c = 3$. It represents the thermodynamic potential for a free colored quark (antiquark) at high temperature, i.e, the color charges are completely ionized and reside at those maxima of Figs. 1 and 5. At low temperature ($T \leq T_s$), the matter part can be written as

$$\Omega_{q(\bar{q})}^{\Phi_0,\bar{\Phi}_0\to 0} = -2N_f T\int\frac{d^3p}{(2\pi)^3}\ln\left(1+e^{-\beta N_c(\epsilon_q\mp\mu)}\right) \quad (16)$$

This represents the thermodynamic potential for a composite color singlet object containing three quarks(antiquarks) in a same color state with the same energy. In other words, the PL dynamically confines three colored quarks in a same energy state, which finally forms a color neutral object like a baryon. This is because $Z(N_c)$ charge gets frozen in color singlet states like baryon as shown in Fig. 7.

Pure gauge sector: The thermodynamic potential, $\Omega_g^{\rm PL}$ in (14) describes the interaction of the spatial gluons with the PL at finite T. The $Z(N_c)$ domains are plentiful (viz., Eq.(10)) as the gluon dynamics are solely governed by PL and its conjugate fields through (8). In asymptotically high temperatures ($T \gg T_c$), $\Phi_0, \bar{\Phi}_0 \to 1$, $A_0 \to 0$, one recovers free gluon gas from $\Omega_g^{\rm PL}$ as

$$\Omega_g^{\Phi_0,\bar{\Phi}_0\to 1} = 2(N_c^2-1)T\int\frac{d^3p}{(2\pi)^3}\ln\left(1-e^{-\beta\epsilon_g}\right) \quad (17)$$

where the spatial gluons are completely ionized. At low



FIG. 8: (a) An exchange of a pair of massive magnetic gluons between two PLs (b) gluon condensates when $\Phi = 0$.

temperature ($T \leq T_s$) and $\Phi_0 = 0$, Ω_g^{PL} becomes

$$\Omega_g^{\Phi_0,\bar{\Phi}_0\to 0} = 2T\int \frac{d^3p}{(2\pi)^3}\left[\ln\left(1-e^{-N_c\beta\epsilon_g}\right)^2 + \ln\left(1-e^{2\pi i/N_c}e^{-\beta\epsilon_g}\right) + \ln\left(1-e^{-2\pi i/N_c}e^{-\beta\epsilon_g}\right)\right] \quad (18)$$

Equation(18) can be viewed in the following way: (i) the first term indicates that the PL confines N_c number of spatial gluons in a same energy state representing a glueball. There are two such copies which is consistent with $SU(3)$ gauge theory as $8\otimes 8\otimes 8$ generates only two singlet glueball states. Obviously, the $Z(N_c)$ charge is frozen in the color singlet glueballs through the localization of $Z(N_c)$ charge in one of the true $Z(N_c)$ vacua (Figs. 1, 2 and 7). (ii) The two spatial gluons in the second and third terms are conjugate to each other but distinguished by $Z(N_c)$ phase. These two conjugate spatial gluons can be visualized in the nonzero Φ domain as an exchange of a pair of massive spatial gluons between two PLs (Fig. 8(a)). This is because Euclidean time reflection does not allow an exchange of a massless pair of spatial gluons [14]. It generates a magnetic screening mass ($\sim g^2T$), a solely non-perturbation correction to the electric screening mass, obtained from the correlation function of the PLs [14]. This in turn provides a non-perturbative mass gap at $T > T_s$ to prevent those excitations of color charges having energy lower than it. In a confined phase ($T \leq T_s$; $\Phi \sim 0$), PL disappears but this two magnetic gluons keep circulating in a loop of virtual color and anti-color charges as a mass dimension four condensate [15] in a non-perturbative vacuum (Fig. 8(b)). These two magnetic gluons should not contribute directly to the thermodynamics in a confined phase. Rather they provide a ground state pressure which is negative and unbound from below which can be viewed as the general confining background of strong interactions.

Polyakov Loop and recombination of partons: When Φ_0 and $\bar{\Phi}_0 \to 0$, i.e, $T \to T_s^+$, the symmetry in $Z(3)$ dictates that the confined phase appears as a color singlet object from the dynamical recombination of three partons as given in (16) and in the first term in (18) plus a vacuum energy density. This would solely describe the thermodynamic properties of color singlet structures like baryon, antibaryon, meson and glueballs. In addition the hybrid structures also appear if one combines the gluonic part with the matter part. This essentially boils down to the fact that the $SU(N_c)$ color singlet restriction supplemented with PL dynamically provides the basis for the recombination of partons for hadronisation from quark-gluon plasma when it cools down to T_s through T_c. It also leads to the scaling law of the valence partons [6, 16] of the eliptic flow of the identified hadrons in heavy-ion collisions, a direct evidence of deconfined phase [17]

Acknowledgment: CAI would like to thank UGC for the financial support. We note that after completion of this work a similar work [18] but in different perspective came to our notice.

---

[1] J. I. Kapusta and C. Gale, *Finite-Temperature Field Theory Principles and Applications*, Second Edition (Cambridge University Press, Cambridge, England, 1996)

[2] H. Weyl, The Classical Groups (Princeton U.P., Princeton, NJ, 1946).

[3] G. Auberson *et al.*, J. Math. Phys. **27**, 1658 (1986).

[4] B. Müller, *The Physics of Quark-Gluon Plasma*, First Edition (Springer-Verlag, Berlin, Germany, 1985); L. Turko, Phys. Lett. B**104**, 153 (1981); H-T Elze and W. Greiner, Phys. Lett. B**179**, 385 (1986); K. Kusaka, Phys. Lett. B**269**, 17 (1991); I. Zakout and C. Greiner, Phys. Rev. C**78**, 034916 (2008); ⟨ arXiv:1107.5497 ⟩.

[5] A. Ansari and M. G. Mustafa; Nucl. Phys. A**539**, 752 (1992); Phys. Rev. C**55**, 2005 (1997); *ibid.* **56**, 420 (1997); M. G. Mustafa, Phys. Lett. B**318**, 517 (1994); Phys. Rev. D**49**, 4634 (1994); M.G. Mustafa, A. Sen, and L. Paria, Eur.Phys. J. C**11** 729 (1999).

[6] R. Abir and M. G. Mustafa, Phys. Rev. C**80**, 051903 (R) (2009).

[7] R. D. Pisarski, Phys. Rev. D**62**, 111501 (2000); A. Dumitru and R. D. Pisarski, Phys. Lett. B**525**, 95 (2002).

[8] F. Fukushima, Phys. Lett. B**591**, 277 (2004); Phys. Rev. D**68**, 045004 (2003); C. Ratti, M. A. Thaller, and W. Weise, Phys. Rev. D**73**, 014019; S. K. Ghosh, T. K. Mukherjee, M. G. Mustafa, and R. Ray, Phys. Rev. D **73**, 114007 (2006); *ibid.* **77**, 094024 (2008); S. Mukherjee, M. G. Mustafa, and R. Ray, *ibid.* **75**, 094015 (2007); C. Sasaki, B. Friman, and K. Redlich, Phys. Rev. D **75** 074013 (2007); A. Bhattacharyya, P. Dev, S. K. Ghosh, and R. Ray, Phys. Rev. D **82**, 014021 (2010).

[9] Y. Nambu and G. Jona-Lasinio, Phys. Rev. **122**, 345 (1961); *ibid* **124**, 246 (1961).

[10] T. Hatsuda and T. Kunihiro, Phys. Rep. **247**, 221 (1994); T. Kunihiro, Phys. Lett. B **271**, 395 (1991).

[11] P. N. Meisinger, M. C. Ogilvie, and T. R. Miller, Phys. Lett. B**585**, 149 (2004).

[12] C. Sasaki and K. Redlich, ⟨arXiv:1204.4330 [hep-ph]⟩.

[13] M. Cheng *et al.*, Phys. Rev. D**77**, 014511 (2008).

[14] P. Arnold and L. G. Yaffe, Phys. Rev. D**52**, 7208 (1995).

[15] P. Chakraborty and M. G. Mustafa, Phys. Rev. D**85**, 056002 (2012).

[16] R. Fries *et al.*, Phys. Rev. Lett. **90**, 202303 (2003); V. Greco, C. M. Ko, and P. Levai, *ibid.* **90**, 202303.

[17] PHENIX Collaboration A. Adare et al., Phys. Rev. Lett. **98**, 162301 (2007).

[18] M. Asakawa, S. A. Bass, and B. Müller, ⟨arXiv:1208.2426 [nucl.th]⟩.